



Predictors of Mothers' Care Seeking Behavior for Common Childhood Illnesses: Findings From the Afghanistan Health Survey 2015



Essa Tawfiq^{1*}, Khwaja Mir Islam Saeed^{2,3}, Sayed Ali Shah Alawi⁴, Jammalluddin Jawaid⁵, Syed Nasir Hashimi⁶

Abstract

Background: Mothers' care seeking behavior for childhood illness is a key factor of utilizing healthcare for children. We examined predictors of mothers' care seeking for common childhood illnesses.

Methods: This was a cross-sectional study, using data from the Afghanistan Health Survey (AHS) 2015. Data were used from women who sought healthcare for their unwell children. The women were asked whether their children were sick with fever, cough, faster breathing, or diarrhea in the past 2 weeks. The outcome variable was defined as whether the mother sought healthcare for her unwell child from a public clinic, a private clinic, or from a pharmacy store. The Andersen's healthcare seeking behavior model was used and multinomial regression analysis applied.

Results: There were 4979 women, aged 15-49 years, whose under-5 children were sick in the past 2 weeks. Thirty-nine percent of women sought healthcare for their children from a health provider. Mother's age, child's age, child's sex, socioeconomic status, mothers' perceived severity of childhood illness, and number of under-5 children were predictors of mothers' care seeking behavior. The likelihood of healthcare seeking was lower for older children (Adjusted odds ratio [OR] [95% CI]: 0.51 [0.37-0.70] from public clinics; 0.33 [0.23-0.47] from private clinics; 0.36 [0.22-0.61] from pharmacy stores), and for girls (Adjusted OR [95% CI]: 0.74 [0.59-0.93] from private clinics). The likelihood of healthcare seeking was greater for children whose mothers knew symptoms of childhood illness (Adjusted OR [95% CI]: 2.97 [1.44-6.16] from public clinics; 7.20 [3.04-17.04] from pharmacy stores). The likelihood of healthcare seeking for children was greater in older mothers (Adjusted OR [95% CI]: 1.54 [1.11-2.12]). It was less likely for the mothers who have more children to seek healthcare for their children (Adjusted OR [95% CI]: 0.53 [0.43-0.65] from public clinics; 0.61 [0.48-0.79] from private clinics; 0.51 [0.37-0.69] from pharmacy stores).

Conclusion: Health policy-makers may opt to use our findings, particularly mothers' knowledge (perceived severity) of symptoms of childhood illness to develop interventions to enhance timely assessment and effective treatment of common preventable childhood illnesses.

Keywords: Care Seeking Behaviour, Childhood Illness, Afghanistan

Copyright: © 2023 The Author(s); Published by Kerman University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Tawfiq E, Saeed KMI, Alawi SAS, Jawaid J, Hashimi SN. Predictors of mothers' care seeking behavior for common childhood illnesses: findings from the Afghanistan health survey 2015. *Int J Health Policy Manag.* 2023;12:7598. doi:10.34172/ijhpm.2023.7598

Article History:

Received: 5 August 2022

Accepted: 28 October 2023

ePublished: 26 November 2023

*Correspondence to:

Essa Tawfiq

Email: e.tawfiq@unsw.edu.au

Background

Child death rates are unacceptably high in low- and middle-income countries (LMICs), and poor healthcare seeking behavior for preventable childhood illnesses is a key obstacle to increasing utilization of available healthcare and reducing morbidity and mortality among children under 5 years of age (under-5 children). Globally, in 2020 five million children died before reaching their fifth birthday.¹ Childhood illnesses such as fever, diarrhea, and acute respiratory infection pose serious health issues in children in LMICs²; acute respiratory infection, diarrhea and/or fever are the cause of death in 3 out of 4 under-5 children who die in LMICs.² In Afghanistan, despite the gradual decline in under-5 mortality rate from 178 deaths per 1000 live births in 1990, to 129 in 2000 and to 58 deaths per 1000 live births in 2020,¹ it is anticipated that the under-5 mortality would rise due to the disruption of and

reduction in the amount of donor funding to the Afghanistan health system since the takeover of political power by the Taliban in August 2021.^{3,4} The pressure on the health system to collapse has been compounded by the ongoing complex humanitarian crises and declining economic status of the Afghan society,^{5,6} which contributed to the growing concern on increasing poverty, hunger, unemployment, migration in the population, and malnutrition, diseases, and deaths in children. The United Nations (UN) and partners launched funding appeals for Afghanistan at several international platforms, and the international donors stepped up to keep Afghanistan from a worsening humanitarian crisis by saving the health system from collapse. With pledges from the donors to continue funding the health system, it is likely that the health system would be able to provide basic healthcare to Afghan residents, even if the funding amounts would not be

Key Messages

Implications for policy makers

- There is a lot of room to improve healthcare seeking for childhood illness, as our findings showed that healthcare was sought for only 2 out of 5 unwell under-5 children. Considering the Afghanistan context, the United Nations (UN) agencies and international donors in the health sector may opt to discuss with the Ministry of Health to explore and adapt culturally appropriate health education interventions, with focus on common childhood illnesses, at the primary care level.
- Substantial gender inequity exists in the utilization of health services for under-5 children. Policy interventions aimed at improving gender equity in utilization of health services for children are desirable.
- Mothers' knowledge of symptoms of childhood illness, which measures perceived severity of childhood illness in our study, is crucial to reduce childhood illnesses and deaths. Our findings provided evidence that the more a mother knows the symptoms of childhood illness the greater would be the likelihood that the mother seeks healthcare for her child.
- Policy-makers may opt to use our findings and develop policies and strategies to improve mothers' care seeking behavior for unwell children, particularly for girls.

Implications for the public

The likelihood of healthcare seeking for younger children is higher than older children. This can be considered a good practice because younger children get more serious illness and need more attention by their parents. On the other hand, our study shows that when a mother has more children, it is less likely she seeks healthcare for her unwell children in comparison to a mother who has fewer children. Furthermore, our study shows that seeking healthcare from private clinics for girls compared to boys is less likely. This may suggest a preferential attitude of parents towards their male children. Parents should pay equal attention to their children and take them to a health provider when the children are sick, regardless of the child's sex and how many children live in the household. Parents, particularly mothers are encouraged to learn more about symptoms of common childhood illness through participating in health education sessions at health facilities or any other platforms suitable in the society.

as much as the UN and partners appealed for Afghanistan.⁷

The Integrated Management of Childhood Illness (IMCI) strategy has been implemented in LMICs since late 1990s.⁸⁻¹⁰ In Afghanistan, significant improvement has been achieved over the past two decades,¹¹ which may have contributed to the implementation of IMCI strategy and subsequent improvement in child health and child well-being.¹² IMCI strategy aims to prevent deaths and diseases while improving the quality of care for under-5 children. It consists of three parts: (i) Improving the skills of healthcare workers by providing training and guidelines, (ii) Improving how healthcare systems are organized and managed, including access to supplies, and (iii) Visiting homes and communities to promote good child rearing practices and good nutrition, while encouraging parents to seek healthcare for their children when the children get sick.

Improving healthcare seeking behavior of mothers can contribute to the timely assessment and effective treatment of common causes of childhood illness, and reducing deaths among under-5 children in LMICs.¹³ However, considerable number of mothers do not take their unwell children to health facilities,¹⁴ due to a number of factors, including household income, mother's illiteracy, mother's age, type of support mothers receive from their husbands, financial hardship, and distance to health facilities.¹⁴⁻¹⁶ Sociocultural factors of parents play a significant role in the decision to seek healthcare for their children's illnesses.¹⁶⁻¹⁹

In LMICs, research shows that mothers usually have little knowledge of common symptoms of childhood illness.^{16,17} Research shows that mothers seek healthcare from traditional healers before opting for health facilities for their children's illnesses.^{16,17,19} Several studies reported that characteristics such as child's age, child's sex, mother's education level, distance to a health facility, residential area, and household socioeconomic status are key influential factors of healthcare

seeking practices for childhood illness.¹⁷⁻²⁶

Timely healthcare seeking by the parents may prevent morbidity and mortality in under-5 children,^{16,27} and yet improving healthcare seeking behavior remains a challenge today.²⁸ Delays in seeking healthcare by mothers to take their children to a health facility may be associated with higher child mortality.²⁹ Delays in seeking healthcare by a mother can lead to the worsening of a child illness which makes the medical care provided to the child to be ineffective. Timely healthcare seeking behavior of parents for their children is a crucial factor in preventing child morbidity.³⁰ In Afghanistan, influential factors associated with healthcare seeking behavior of mothers for their children have remained insufficiently studied.

The aim of this study was to examine predictors of mothers' care seeking behavior for common childhood illnesses. The findings from this study may have the potential to influence healthcare policies and interventions to encourage timely healthcare seeking behavior and improve diagnoses and effective treatments of childhood illness.

Methods

Study Design

The study used secondary data from the Afghanistan Health Survey 2015 (AHS 2015). This was a cross-sectional study, using data collected from households for the AHS 2015.³¹ Data were collected between July and December 2015 from 23 137 households across all 34 provinces of Afghanistan. In this study we included ever-married women, aged 15-49, who had under-5 children. As part of the AHS 2015 questionnaires, women were asked whether they sought healthcare from a health provider for their under-5 children who were sick because of fever, cough, diarrhea, or faster breathing in the past 2 weeks prior to the survey. A total of 24 427 ever-married women, aged 15-49 years, who had under-5 children were

interviewed, and 5532 of them answered that their children were sick in the past two weeks prior to the survey. A total of 4979 women with complete information who met the above criteria were included for data analysis in this study (Figure 1).

Sample Design and Data Collection

The AHS 2015 used a stratified two-stage sample design.³¹ Within each province (stratum), 30 clusters were randomly selected from the list of clusters provided by the central statistics office. Within each cluster, 23 households were selected randomly by the field teams.³¹ The field teams were trained surveyors and were recruited from provinces to minimize selection bias during data collection. Selection bias could have happened if the surveyors were not from the respective provinces. In such scenarios, the surveyors may have dropped the sampled cluster(s) due to security concerns or logistic challenges in the area.

During the survey, two types of questionnaires were used: the household questionnaire, and the women's and children's questionnaire. The household questionnaire was used to identify women who were eligible to be interviewed for the questions in the women's and children's questionnaire. During the interview, the women who had under-5 children were asked the following questions related to acute childhood illness.

1. In the past 2 weeks, has (name of the child) had diarrhea?
2. In the past 2 weeks, has (name of the child) been ill with a fever?
3. In the past 2 weeks, has (name of the child) been ill with a cough?
4. In the past 2 weeks, has (name of the child) had fast, or short and rapid breathing, or difficulty breathing?

If the response was "yes" for any of the above questions, the following question was asked.

5. Did you seek any advice or treatment from any source?
If the response was "yes" the following question was asked.
6. From where you FIRST sought advice or treatment?

We used a modified version of the Andersen's behavioral model for health services that was employed in previous studies,^{32,33} and created a theoretical framework for mothers' care seeking behavior for unwell children (Figure 2).

Statistical Analysis

Data from women whose under-5 children were sick with fever, diarrhea, cough, or had rapid breathing or difficulty breathing in the past 2 weeks were included and analyzed. Cases with complete information on the outcome variable and explanatory variables were included for data analysis. The




Figure 1. Number of Mothers Entered in the Study.

outcome variable (a four-categorical variable) was created as healthcare was sought from a public clinic (category=1), from a private clinic (category=2), from a pharmacy store (category=3) vs. healthcare was not sought (category=0), using the questions "Did you seek any advice or treatment from any source?" and "From where you first sought advice or treatment?"

The explanatory variables were mother's age (15-29 years, 30-39 years, and 40-49 years), child's age (<2 years, 2 to under 4 years, and 4 to under 5 years), child's sex, mother's education level (no formal education, completed primary education, completed secondary or higher education), socioeconomic status (quintile), number of under-5 children in the household (1 child, 2-3 children, and ≥4 children), access to radio (never listened to radio vs. listened to radio), access to TV (never watched TV vs. watched TV), and mother's knowledge of symptoms of childhood illness (knew no symptom, knew 1 symptom, knew 2-3 symptoms, knew 4-7 symptoms). We used the mothers' knowledge of symptoms as a measure of severity of childhood illness perceived by the mother (See Table 1). The level of perceived severity was determined by the number of symptom(s) a mother knew.

The variable on socioeconomic status was created using the principal component analysis, using the household ownership of assets, household use of amenities of life, and construction material used for the household properties. Using the "pca" function in Stata version 13, the correlation matrix between the variables of household assets, household use of amenities, and construction material used for the household properties




Figure 2. Theoretical framework for Mothers' Healthcare Seeking Behavior for Unwell Children.

Table 1. Women's Perceived Severity of Childhood Illness

The Question on Severity of Childhood Illness	Response Options (Symptoms)
What type of symptoms would cause you to take your child to a health facility right away?	1. Child cannot suckle breastmilk
	2. Child becomes sicker
	3. Child develops a fever
	4. Child has fast breathing
	5. Child has difficulty breathing
	6. Child has blood in stool
	7. Child is drinking poorly
	8. Others (specify)
Instructions to surveyor(s). Record all responses the respondent provides. Do not make any suggestion.	

was obtained. The first component, which explained 11% variance in the original variables related to household assets, amenities, and construction material, was used to create quintile of socioeconomic status.

Multicollinearity between explanatory variables was examined by using the variance inflation factors (VIF) function in Stata version 13, and we found that the highest VIF value was 1.29 which was for the variable on socioeconomic status. Given the highest VIF value was 1.29, we concluded that there was no indication of substantial multicollinearity between variables.³⁴

For multivariable analysis, a generalized linear model with a multinomial outcome was specified as shown below.

$$Y_{ij} = \beta_0 + \sum_{k=1}^K \beta_j X_{ij} + \varepsilon_{ij}$$

Y_{ij} refers to the outcome variable for woman i (whether she sought healthcare from a public clinic; whether she sought healthcare from a private clinic, whether she sought healthcare from a pharmacy store, with the response of “yes/no” for each of them), with j category of explanatory variables. X_{ij} denotes a vector of explanatory variables, and k refers to the number of explanatory variables. β_0 stands for the intercept term, and the exponential of β_j provides the odds ratio (OR) of seeking healthcare for each category of explanatory variables. ε_{ij} refers to the error term. Provincial weights based on number of households were used. Data were analyzed in Stata version 13 using the “svyset,” “strata,” and “svy” functions before running the “mlogit” function in the multinomial logit model to account for clustering and weighting. The ORs were obtained by the “mlogit, rrr” function.

Results

Table 2 shows that there were 4979 ever-married women, aged 15-49 years, whose under-5 children were unwell in the past 2 weeks. Thirty-nine percent of women sought healthcare from a health provider for their children who had fever, cough, faster breathing, or diarrhea. There were significant differences between the mothers who sought healthcare and those who did not seek healthcare for their children. Higher proportions of older mothers sought healthcare for their children, and for a higher proportion of children <2 years healthcare was sought. For a significantly higher proportion of boys, and for a significantly higher proportion of children

whose mothers knew ≥ 4 symptoms of childhood illness healthcare was sought. In households with 1 child, for a substantially higher proportion of children healthcare was sought. An unusual finding was that for a higher proportion of children whose mothers watched TV, healthcare was not sought (details in Table 2). Weighted proportions for the baseline characteristics of mothers and children are provided in Supplementary file 1.

Figures 3A and 3B show the proportions of mothers who sought healthcare for their children, by health provider type, and the proportions of children for whom healthcare was sought, by illness type, respectively. Public clinics were the facilities from where healthcare for children was sought by 46.1% of mothers, followed by private clinics (39.7%), and pharmacy stores (14.2%). The most common illnesses the children had, and healthcare was sought for, were diarrhea (63.8%), followed by fever (60.2%), cough (54.6%), and faster breathing (38.6%).

Table 3 shows results from multivariable analysis. Mother's age, child's age, child's sex, socioeconomic status, mothers' perceived severity of childhood illness, and number of under-5 children were predictors of mothers' care seeking behaviour for childhood illness. It shows that child's age is a significant predictor of mothers' care seeking behavior for childhood illness, as the likelihood of mothers' care seeking was significantly low for older children (Adjusted OR 0.53 [0.41-0.69] from private clinics; 0.63 [0.46-0.88] from pharmacy stores for children 2 years to under-4 years), and [Adjusted OR 0.51 [0.37-0.70] from public clinics, 0.33 [0.23-0.49] from private clinics, and 0.36 [0.22-0.61] from pharmacy stores for children 4 years to under-5 years), compared to under-2 years old. It was more likely for mothers, aged 40-49 years, to seek healthcare for children (Adjusted OR 1.54 [1.11-2.12] from private clinics) compared to mothers, aged 15-29 years (details in Table 3).

Data on the reasons for not seeking healthcare were available for and analyzed from 613 out of the 3032 mothers who did not seek healthcare for their unwell children. Figure 4 shows that 31.5% of mothers thought it was unnecessary to seek healthcare, 17.5% thought services were expensive, 17.5% responded the clinic was too far, 7.3% thought transportation cost was too expensive, 7.3% expressed security concern, and 7.2% responded that staff in the clinic were unfriendly. Ranging from 1.6% to 2.9% of mothers reported the following

Table 2. Baseline Characteristics of Mothers and Their Under-5 Children

		All women and Children No. (%)	Those Who Sought Care No. (%)	Those Who Did not Seek Care No. (R)	P Value
Total	Women or children	4979 (100.0)	1947 (39.1)	3032 (60.9)	<.001
Age of mother (y)	15-29	2678 (54.0)	985 (50.6)	1693 (55.8)	.001
	30-39	1642 (33.0)	675 (34.7)	967 (31.9)	
	40-49	659 (13.0)	287 (14.7)	372 (12.3)	
Age of child (y)	<2	1782 (36.0)	827 (42.5)	955 (31.5)	<.001
	2-<4	2196 (44.0)	843 (43.3)	1353 (44.6)	
	4-<5	1001 (20.0)	277 (14.2)	724 (23.9)	
Gender of child	Boy	2659 (53.4)	1083 (55.6)	1576 (52.0)	.012
	Girl	2320 (46.6)	864 (44.4)	1456 (48.0)	
Education Level of women	No education	4089 (82.1)	1602 (82.3)	2487 (82.0)	.65
	Primary education	647 (13.0)	245 (12.6)	402 (13.3)	
	Secondary or higher education	243 (4.9)	100 (5.1)	143 (4.7)	
Socioeconomic status	Lowest (quintile)	1007 (20.2)	393 (20.2)	614 (20.3)	.010
	Low	994 (20.0)	412 (21.2)	582 (19.2)	
	Middle	1017 (20.4)	417 (21.4)	600 (19.8)	
	High	984 (19.8)	338 (17.4)	646 (21.3)	
	Highest (quintile)	977 (19.6)	387 (19.9)	590 (19.5)	
Perceived severity of childhood illness	Knowing no symptom	210 (4.2)	53 (2.7)	157 (5.2)	<.001
	Knowing 1 symptom	840 (16.9)	305 (15.7)	535 (17.7)	
	Knowing 2-3 symptoms	3036 (61.0)	1182 (60.7)	1854 (61.2)	
	Knowing ≥4 symptoms	893 (17.9)	407 (20.9)	486 (16.0)	
Number of under-5 children in household	1 child	2138 (42.9)	1051 (54.0)	1087 (35.8)	<.001
	2-3 children	2707 (54.4)	877 (45.0)	1830 (60.4)	
	≥ 4 children	134 (2.7)	19 (1.0)	115 (3.8)	
Women listen to radio	Yes	2739 (55.0)	1040 (53.4)	1699 (56.0)	.07
	No	2240 (45.0)	907 (46.6)	1333 (44.0)	
Women watch TV	Yes	2433 (48.9)	917 (47.1)	1516 (50.0)	.046
	No	2546 (51.1)	1030 (52.9)	1516 (50.0)	

Note: *P* values were obtained from chi-square test for cross-tabulations of characteristics between mothers who sought healthcare and mothers who did not. A two-sample test of proportions was conducted to obtain *P* value for women who sought care (39.1%) and women who did not seek care for their children (60.9%).

reasons for not seeking healthcare for their unwell children: unavailability of female staff in the clinic, inconvenience service hours, religious reasons, and lack of an accompanying man.

Discussion

The discussion has been structured considering the Andersen's model used for the theoretical framework in this study. This study revealed that only 39.1% of women sought healthcare for childhood illness. Previous studies from LMICs reported that 45% to 65% of mothers sought healthcare for childhood illness.^{22,35-37} With respect to predisposing factors, our study showed that child's age is a strong predictor of mothers' care seeking behavior for childhood illness, and our finding supports those previously reported from LMICs.^{22,32,37} We observed lower odds of healthcare seeking for girls, which is consistent with findings from several LMICs.^{32,38-40} A plausible reason could be related to the parents preference for male

children by some cultures, including those in Afghanistan.^{39,41} In our study it was found that mother's age is a strong predictor of mothers' care seeking behavior for childhood illness. A recent study in LMICs showed that mother's age is a potential, but not significant, predictor of mothers' care seeking behavior for childhood illness.³⁵ A plausible reason for our finding could be attributed to the experience mothers gain for seeking healthcare for their children throughout their life span. Our findings that mother's education is not a strong predictor of healthcare seeking for sick children contradicts findings previously reported from LMICs.^{17,32,35} Despite our findings on the effect of mother's education on healthcare seeking behavior, we cannot rule out the pivotal role of mother's education on healthcare seeking for childhood illness because of the abundance of evidence reported from several studies.^{17,32,35} However, our finding that number of children in a household is a strong predictor of mothers' care seeking behavior for childhood illness is consistent with those




Figure 3. (A) Proportions of Maternal Care-Seeking for Children by Health Provider Type and (B) Proportions of Children Receiving Care by Illness Type.

previously reported.³⁵

With respect to enabling factors, our study identified that socioeconomic status is a significant predictor of mothers' care seeking behavior for childhood illness, and this is consistent with previous findings from LMICs.^{22,32,35,42} Our findings that odds of healthcare seeking from private clinics was higher for children from wealthier families support those previously reported.³² Unlike above, our findings that mothers' exposure to radio and television did not affect mothers' care seeking behavior for childhood illness contradict those from previous studies.^{35,43} Although, our findings contradict the significant role of mothers' access to media on care seeking behavior, recent data from Afghanistan indicate the significant effects of women's exposure to media on care seeking behavior.^{44,45}

With respect to perceived severity of illness, which could prompt the need to seek healthcare, our findings support those

previously reported from LMICs.^{22,32,33} Unlike these studies, however, our findings provided a dose-response relationship between the number of symptoms of childhood illness that a mother knows and the likelihood of healthcare seeking for the unwell children. This means the more symptoms a mother knows the higher would be the likelihood of healthcare seeking for the unwell children. A previous study found that mothers' awareness of childhood illness, perceived importance of early treatment, and perceived severity were significant predictors of mothers' care seeking behavior for childhood illness.²² The authors defined perceived illness severity when a mother thought that her sick child was severely ill. Another study found strong associations between mothers' knowledge of symptoms in a child and healthcare seeking for the sick child,³² and examined the symptoms related to diarrhoea in children who received health services for the treatment of




Figure 4. Reasons for not Seeking Healthcare for Unwell Children. Note: 2419 women (out of 3032) did not answer this question.

Table 3. Predictors of Mothers' Care Seeking for Childhood Illness by Health Facility Type (N = 4979)

		Public Clinics	Private Clinics	Pharmacy Stores
		Adjusted OR (95% CI)	Adjusted OR (95% CI)	Adjusted OR (95% CI)
Age of mother (y)	15-29	Ref	Ref	Ref
	30-39	1.21 (0.97-1.52)	1.14 (0.88-1.48)	1.16 (0.83-1.63)
	40-49	1.03 (0.78-1.38)	1.54 ^b (1.11-2.12)	1.32 (0.81-2.13)
Age of child (y)	<2	Ref	Ref	Ref
	2 to <4	0.81 (0.64-1.01)	0.53 ^c (0.41-0.69)	0.63 ^b (0.46-0.88)
	4 to <5	0.51 ^c (0.37-0.70)	0.33 ^c (0.23-0.47)	0.36 ^c (0.22-0.61)
Gender of child	Boy	Ref	Ref	Ref
	Girl	0.95 (0.78-1.16)	0.74 ^b (0.59-0.93)	1.15 (0.85-1.56)
Education level of woman	No education	Ref	Ref	Ref
	Primary education	0.88 (0.64-1.22)	0.90 (0.62-1.31)	0.76 (0.49-1.18)
	Secondary education	1.58 (0.94-2.67)	0.94 (0.57-1.52)	1.23 (0.63-2.42)
Socioeconomic status	Lowest (quintile)	Ref	Ref	Ref
	Low	1.21 (0.94-1.57)	1.20 (0.86-1.67)	1.42 (0.87-2.32)
	Middle	1.03 (0.78-1.35)	1.20 (0.87-1.64)	1.12 (0.67-1.89)
	High	0.73 (0.53-1.01)	1.01 (0.71-1.42)	1.20 (0.70-2.04)
	Highest (quintile)	0.39 ^c (0.26-0.59)	1.72 ^c (1.24-2.40)	1.49 (0.90-2.46)
Perceived severity of childhood illness	Knows none	Ref	Ref	Ref
	Knows 1 symptom	1.67 (0.79-3.54)	1.37 (0.75-2.51)	2.01 (0.81-5.01)
	Knows 2-3 symptoms	2.29 ^a (1.13-4.65)	1.40 (0.82-2.38)	3.98 ^c (1.71-9.28)
	Knows ≥ 4 symptoms	2.97 ^c (1.44-6.16)	1.67 (0.94-2.97)	7.20 ^c (3.04-17.04)
Number of under-5 children in household	1 child	Ref	Ref	Ref
	2-3 children	0.53 ^c (0.43-0.65)	0.61 ^c (0.48-0.79)	0.51 ^c (0.37-0.69)
	≥4 children	0.14 ^c (0.06-0.35)	0.27 ^c (0.13-0.56)	0.29 (0.04-2.08)
Listens to radio	No	Ref	Ref	Ref
	Yes	1.05 (0.85-1.30)	1.11 (0.90-1.38)	1.11 (0.82-1.50)
Watches TV	No	Ref	Ref	Ref
	Yes	0.96 (0.76-1.22)	0.98 (0.80-1.21)	1.30 (0.91-1.84)

Abbreviations: OR, odds ratio; CI, confidence interval.

^a $P < .05$, ^b $P < .01$, ^c $P < .001$.

diarrhoea, and the symptoms related to cough and fever in children who received health services for the treatment of cough and fever. In our study, we defined mothers' perceived severity of childhood illness when a mother knew symptoms in a sick child that alerted her to take the child to a health provider right away. Symptoms such as inability of a child to suckle breastmilk, become sicker, develop a fever, have fast breathing, have difficulty in breathing, have blood in stool, or drink poorly were considered mothers' perceived severity of childhood illness; likewise, the number of symptoms indicated the degree of illness severity. In our study, there were few mothers who provided responses which were not among the seven symptoms we used as the mothers' perceived severity of illness. These responses were heterogenous and it was not possible to categorize them in a specific category, and they were not analysed.

Our findings on mothers' perceived severity of childhood illness, predictors of mothers' care seeking behavior, and the low prevalence of healthcare seeking behavior by mothers for childhood illness have the potential to influence health policies

and interventions to promote healthcare seeking behavior for childhood illness. This is particularly important for the current situation to prioritize more targeted interventions to maximize the use of limited resources available in the health sector in Afghanistan. Health interventions which accommodate activities to improve mothers' care seeking behavior for childhood illness may lead to the rationale use of available limited resources at the primary care level. This may help reduce the utilization of more expensive therapeutic interventions at the secondary care level.

The simple question on mothers' knowledge of symptoms of childhood illness with related response options in our study can be used to target health education activities related to common childhood illnesses. Research shows that early healthcare seeking for childhood illness can affect the use of health services offered from health facilities at the primary care level versus at the secondary care level.³⁷ Our findings showed that one third of women, who did not seek healthcare for their unwell children, thought that care was not necessary. This emphasizes the need to focus more on educating women

on symptoms that prompt healthcare seeking for unwell children. This should be noted that these findings are from data collected in 2015, which means that the extent of the reasons for not seeking healthcare may have changed over time, especially after the take-over of political power by the Taliban in August 2021. Under the Taliban's rule, more women may not seek healthcare for their children because of the higher cost of transportation, services, and because of the shortage of essential drugs and health staff, even though security concern may have decreased in most parts of Afghanistan.

The strengths of our study are the use of data from a nationally representative database, and the use of questionnaires and having a study design similar to those used in the demographic health surveys.³⁵ This allows generalization of our findings at the national level, and the comparison of our findings with findings from other LMICs. The sample size is sufficiently large, and this can increase the validity of our findings.

The limitations of our study are the chance of social desirability and recall bias,⁴⁶ though the change of recall bias may have been minimal because the interviewee mother was asked about the events for her child in the past 2 weeks prior to the survey. Another limitation is the lack of data on access to health facilities. Research shows that there is a strong association between distance to health providers and healthcare seeking behavior for childhood illness.^{40,43,47} A further limitation of our study is that the variable used to measure socioeconomic status did not capture it very well.

Future research may examine the association between distance to health facilities and healthcare seeking behavior, including delays in seeking healthcare and use of health services from primary care facilities versus hospitals. Another venue to explore may be the cost of using health services, including transportation cost and its association with healthcare seeking for childhood illness.

Conclusion

Effective health policy interventions are needed to enhance mothers' care seeking behavior for childhood illness. Our findings reinforce the need for targeted interventions to enhance mothers' care seeking behavior for common childhood illnesses. The targeted interventions should focus on improving mothers' perceived severity of childhood illness through increasing their knowledge of symptoms that would prompt actions to take the unwell child to a health provider. The interventions should focus to encourage parents to seek healthcare for their under-5 children, particularly those <2 years old, regardless of the child's sex and number of children living in the household.

Acknowledgements

We thank the Silk Route Training and Research Organization (SRTRO) for sharing the data the organization collected for the AHS 2015.

Ethical issues

Secondary data were used, and it was not required to obtain an ethics approval. The data used in this study did not include identifiable personal information.

Competing interests

Authors declare that they have no competing interests.

Authors' contributions

Conceptualization: Essa Tawfiq, Khwaja Mir Islam Saeed, and Sayed Ali Shah Alawi.

Data curation: Essa Tawfiq and Khwaja Mir Islam Saeed.

Methodology: Essa Tawfiq and Khwaja Mir Islam Saeed.

Formal analysis: Essa Tawfiq.

Visualization: Essa Tawfiq and Khwaja Mir Islam Saeed.

Writing—original draft: Essa Tawfiq.

Writing—review & editing: Essa Tawfiq, Khwaja Mir Islam Saeed, Sayed Ali Shah Alawi, Jammalluddin Jawaid, and Syed Nasir Hashimi.

Authors' affiliations

¹The Kirby Institute, University of New South Wales, Sydney, NSW, Australia.

²Global Health Development/Eastern Mediterranean Public Health Network (GHD/EMPHNET), Amman, Jordan.

³Afghanistan Field Epidemiology Training Program, Afghanistan National Public Health Institute, Kabul, Afghanistan.

⁴Afghanistan Pediatric Association, Kabul, Afghanistan.

⁵Agency for Assistance and Development of Afghanistan, Kabul, Afghanistan.

⁶Jhpiego, Kabul, Afghanistan.

Supplementary files

Supplementary file 1. Baseline Characteristics of Women and Their Children.

References

- United Nations Inter-Agency Group for Child Mortality Estimation (UN IGME), Report 2019. Levels & Trends in Child Mortality. UNICEF, 2021. <https://childmortality.org/wp-content/uploads/2021/12/UNICEF-2021-Child-Mortality-Report.pdf>. Accessed June 10, 2022.
- Kebede ZT, Gelaye KA, Merid MW, et al. High common childhood illnesses while treatment seeking behavior of mothers'/care givers' was low at rural northwest Ethiopia. *Res Sq* [Preprint]. August 7, 2020. Available from: <https://assets.researchsquare.com/files/rs-7641/v3/fa1b4721-fc80-412d-981f-6e7fd73bbd13.pdf?c=1631850080>.
- Britten S, Amer M. Preventing collapse of the Afghan health service. *Lancet*. 2021;398(10310):1481. doi:10.1016/s0140-6736(21)02234-0
- Walraven G, Yousofzai Y, Mirzazada S. The World Bank's health funding in Afghanistan. *Lancet*. 2021;398(10306):1128. doi:10.1016/s0140-6736(21)02015-8
- Afghanistan Situation Supplementary Appeal, UNHCR, The UN Refugee Agency, July – December 2021. <https://reporting.unhcr.org/sites/default/files/2021-12/Afghanistan%2520appeal.pdf>. Accessed June 10, 2022.
- Shah J, Essar MY, Qaderi S, et al. Respiratory health and critical care concerns in Afghanistan. *Lancet Respir Med*. 2022;10(3):229-231. doi:10.1016/s2213-2600(21)00583-x
- Afghanistan: UN Launches Largest Single Country Aid Appeal Ever - Media Release on 11 January 2022. <https://news.un.org/en/story/2022/01/1109492#:~:text=The%20UN%20and%20partners%20launched,requiring%20help%20beyond%20its%20borders>. Accessed June 25, 2022.
- Kolstad PR, Burnham G, Kalter HD, Kenya-Mugisha N, Black RE. The integrated management of childhood illness in western Uganda. *Bull World Health Organ*. 1997;75(Suppl 1):77-85.
- Centers for Disease Control and Prevention (CDC). Health worker performance after training in integrated management of childhood illness—Western province, Kenya, 1996-1997. *MMWR Morb Mortal Wkly Rep*. 1998;47(46):998-1001.
- Simoes EA, Desta T, Tessema T, Gerbresellassie T, Dagnaw M, Gove S. Performance of health workers after training in integrated management of childhood illness in Gondar, Ethiopia. *Bull World Health Organ*. 1997;75(Suppl 1):43-53.
- Tawfiq E, Alawi SAS, Natiq K. Effects of training health workers in integrated management of childhood illness on quality of care for under-5 children in primary healthcare facilities in Afghanistan. *Int J Health Policy Manag*. 2020;9(1):17-26. doi:10.15171/ijhpm.2019.69
- Afghanistan Demographic and Health Survey 2015. Ministry of Public Health, Kabul Afghanistan, and The DHS Program ICF Rockville, Maryland, USA; January 2017.
- Khalid M, Kumari R, Mohan U, Manar M, Singh V. Morbidity profile of preschool children from below poverty line families of Lucknow district, North India. *Int J Adv Res*. 2014;2(11):627-634.
- World Health Organization (WHO). *State of Inequality: Reproductive*,

- Maternal, Newborn and Child Health*. Geneva: WHO; 2015. http://apps.who.int/iris/bitstream/10665/164590/1/9789241564908_eng.pdf. Accessed June 10, 2022.
15. Awoke W. Prevalence of childhood illness and mothers'/caregivers' care seeking behavior in Bahir Dar, Ethiopia: a descriptive community based cross sectional study. *Open J Prev Med*. 2013;3(2):155-159. doi:10.4236/ojpm.2013.32020
 16. Ajibade B, Amoo P, Adeleke M, Oyadiran G, Kolade O, Olagunju RJ. Determinants of mothers health seeking behaviour for their children in a Nigerian teaching hospital. *IOSR J Nurs Health Sci*. 2013;1(6):9-16.
 17. Abegaz NT, Berhe H, Gebretekle GB. Mothers/caregivers healthcare seeking behavior towards childhood illness in selected health centers in Addis Ababa, Ethiopia: a facility-based cross-sectional study. *BMC Pediatr*. 2019;19(1):220. doi:10.1186/s12887-019-1588-2
 18. Tinuade O, Iyabo RA, Durotoye O. Health-care-seeking behaviour for childhood illnesses in a resource-poor setting. *J Paediatr Child Health*. 2010;46(5):238-242. doi:10.1111/j.1440-1754.2009.01677.x
 19. Oberoi S, Chaudhary N, Patnaik S, Singh A. Understanding health seeking behavior. *J Family Med Prim Care*. 2016;5(2):463-464. doi:10.4103/2249-4863.192376
 20. Bennett A, Eisele T, Keating J, Yukich J. *Global Trends in Care Seeking and Access to Diagnosis and Treatment of Childhood Illnesses*. DHS Working Papers No. 116. Rockville, MD: ICF International; 2015.
 21. Lungu EA, Darker C, Biesma R. Determinants of healthcare seeking for childhood illnesses among caregivers of under-five children in urban slums in Malawi: a population-based cross-sectional study. *BMC Pediatr*. 2020;20(1):20. doi:10.1186/s12887-020-1913-9
 22. Simieneh MM, Mengistu MY, Gelagay AA, Gebeyehu MT. Mothers' health care seeking behavior and associated factors for common childhood illnesses, Northwest Ethiopia: community based cross-sectional study. *BMC Health Serv Res*. 2019;19(1):59. doi:10.1186/s12913-019-3897-4
 23. Wambui WM, Kimani S, Odhiambo E. Determinants of health seeking behavior among caregivers of infants admitted with acute childhood illnesses at Kenyatta National Hospital, Nairobi, Kenya. *Int J Pediatr*. 2018;2018:5190287. doi:10.1155/2018/5190287
 24. Adane M, Mengistie B, Mulat W, Kloos H, Medhin G. Utilization of health facilities and predictors of health-seeking behavior for under-five children with acute diarrhea in slums of Addis Ababa, Ethiopia: a community-based cross-sectional study. *J Health Popul Nutr*. 2017;36(1):9. doi:10.1186/s41043-017-0085-1
 25. Ayalneh AA, Fetene DM, Lee TJ. Inequalities in health care utilization for common childhood illnesses in Ethiopia: evidence from the 2011 Ethiopian Demographic and Health Survey. *Int J Equity Health*. 2017;16(1):67. doi:10.1186/s12939-017-0561-7
 26. Bedford KJ, Sharkey AB. Local barriers and solutions to improve care-seeking for childhood pneumonia, diarrhoea and malaria in Kenya, Nigeria and Niger: a qualitative study. *PLoS One*. 2014;9(6):e100038. doi:10.1371/journal.pone.0100038
 27. Olenja J. Health seeking behaviour in context. *East Afr Med J*. 2003; 80(2):61-62. doi:10.4314/eamj.v80i2.8689
 28. Dagnaw AB, Tewabe T, Murugan R. Level of modern health care seeking behaviors among mothers having under five children in Dangila town, North West Ethiopia, 2016: a cross sectional study. *Ital J Pediatr*. 2018; 44(1):61. doi:10.1186/s13052-018-0503-z
 29. Webair HH, Bin-Gouth AS. Factors affecting health seeking behavior for common childhood illnesses in Yemen. *Patient Prefer Adherence*. 2013; 7:1129-1138. doi:10.2147/ppa.s51124
 30. Geldsetzer P, Williams TC, Kirolos A, et al. The recognition of and care seeking behaviour for childhood illness in developing countries: a systematic review. *PLoS One*. 2014;9(4):e93427. doi:10.1371/journal.pone.0093427
 31. Afghanistan Health Survey 2015, Final Report – August 2016. KIT The Royal Tropical Institute, Silk Route Training & Research Organization. <https://www.kit.nl/wp-content/uploads/2018/10/08-23-2016-Afghanistan-Health-Survey-2015-Final-Report.pdf>. Accessed June 25, 2022.
 32. Sreeramareddy CT, Sathyanarayana TN, Kumar HN. Utilization of health care services for childhood morbidity and associated factors in India: a national cross-sectional household survey. *PLoS One*. 2012;7(12):e51904. doi:10.1371/journal.pone.0051904
 33. Sreeramareddy CT, Shankar RP, Sreekumaran BV, Subba SH, Joshi HS, Ramachandran U. Care seeking behaviour for childhood illness-a questionnaire survey in western Nepal. *BMC Int Health Hum Rights*. 2006;6:7. doi:10.1186/1472-698x-6-7
 34. O'Brien RM. A caution regarding rules of thumb for variance inflation factors. *Qual Quant*. 2007;41(5):673-690. doi:10.1007/s11135-006-9018-6
 35. Yaya S, Odusina EK, Adjei NK. Health care seeking behaviour for children with acute childhood illnesses and its relating factors in sub-Saharan Africa: evidence from 24 countries. *Trop Med Health*. 2021;49(1):95. doi:10.1186/s41182-021-00385-1
 36. Breiman RF, Olack B, Shultz A, et al. Healthcare-use for major infectious disease syndromes in an informal settlement in Nairobi, Kenya. *J Health Popul Nutr*. 2011;29(2):123-133. doi:10.3329/jhpn.v29i2.7854
 37. Shimelis T, Schierhout G, Tadesse BT, et al. Timely health care seeking and first source of care for acute febrile illness in children in Hawassa, southern Ethiopia. *PLoS One*. 2022;17(6):e0269725. doi:10.1371/journal.pone.0269725
 38. Degefa G, Gebresslassie M, Meles KG, Jackson R. Determinants of delay in timely treatment seeking for diarrheal diseases among mothers with under-five children in central Ethiopia: a case control study. *PLoS One*. 2018;13(3):e0193035. doi:10.1371/journal.pone.0193035
 39. Khera R, Jain S, Lodha R, Ramakrishnan S. Gender bias in child care and child health: global patterns. *Arch Dis Child*. 2014;99(4):369-374. doi:10.1136/archdischild-2013-303889
 40. Nega B, Bogale KA, Nigussie ZM. Health care seeking behavior and associated factor among mothers/caregivers of under-five children with acute diarrhea in Dangila zuria Woreda, North West Ethiopia. bioRxiv [Preprint]. June 11, 2019. Available from: <https://www.biorxiv.org/content/10.1101/667923v1>.
 41. Odusina EK, Ayotunde T, Kunnuji M, Ononokpono DN, Bishwajit G, Yaya S. Fertility preferences among couples in Nigeria: a cross sectional study. *Reprod Health*. 2020;17(1):92. doi:10.1186/s12978-020-00940-9
 42. Adedokun ST, Adekanmbi VT, Uthman OA, Lifford RJ. Contextual factors associated with health care service utilization for children with acute childhood illnesses in Nigeria. *PLoS One*. 2017;12(3):e0173578. doi:10.1371/journal.pone.0173578
 43. Budu E, Seidu AA, Ameyaw EK, et al. Factors associated with healthcare seeking for childhood illnesses among mothers of children under five in Chad. *PLoS One*. 2021;16(8):e0254885. doi:10.1371/journal.pone.0254885
 44. Tawfiq E, Azimi MD, Feroz A, et al. Predicting maternal healthcare seeking behaviour in Afghanistan: exploring sociodemographic factors and women's knowledge of severity of illness. *BMC Pregnancy Childbirth*. 2023;23(1):561. doi:10.1186/s12884-023-05750-y
 45. Tawfiq E, Fazli MR, Wasiq AW, Stanikzai MH, Mansouri A, Saeedzai SA. Sociodemographic predictors of initiating antenatal care visits by pregnant women during first trimester of pregnancy: findings from the Afghanistan Health Survey 2018. *Int J Womens Health*. 2023;15:475-485. doi:10.2147/ijwh.s399544
 46. Latkin CA, Edwards C, Davey-Rothwell MA, Tobin KE. The relationship between social desirability bias and self-reports of health, substance use, and social network factors among urban substance users in Baltimore, Maryland. *Addict Behav*. 2017;73:133-136. doi:10.1016/j.addbeh.2017.05.005
 47. Sisay S, Endalew G, Hadgu G. Assessment of mothers/care givers health care seeking behavior for childhood illness in rural Ensaro district, North Shoa zone, Amhara region, Ethiopia 2014. *Glob J Life Sci Biol Res*. 2015;1(1):15.